



ISSN: 2321 9114

AJEONP 2013; 1 (2): 38-40

© 2013 AkiNik Publications

Received 03-10-2013

Accepted: 15-11-2013

**Sarah L. Miller**

Department of Chemistry, University  
of Alabama in Huntsville  
Huntsville, AL 35899, USA

**William A. Haber**

Missouri Botanical Garden  
St. Louis, MO 63166, USA  
Apdo. 50-5655, Monteverde  
Puntarenas, Costa Rica, Central  
America

**William N. Setzer**

Department of Chemistry, University  
of Alabama in Huntsville  
Huntsville, AL 35899, USA

## Chemical composition of the leaf essential oil of an undescribed species of *Zanthoxylum* from Monteverde, Costa Rica

Sarah L. Miller, William A. Haber, William N. Setzer

### Abstract

The leaf essential oil of *Zanthoxylum* sp. A “Peñas Blancas” (Rutaceae), from Monteverde, Costa Rica, was isolated by hydrodistillation and analyzed by GC-MS. A total of 12 compounds were identified in the leaf oil accounting for 100% of the total composition. The most abundant components of the essential oil of *Zanthoxylum* sp. A “Peñas Blancas” were citronellal (23.3%), limonene (16.7%), *trans*-pinocamphone (11.6%), and  $\beta$ -pinene (10.8%). A cluster analysis comparison with previously published leaf oil compositions of *Zanthoxylum* species revealed a high degree of interspecific as well as intraspecific variation in the genus.

**Keywords:** *Zanthoxylum*; composition; citronellal; limonene; *trans*-pinocamphone;  $\beta$ -pinene; cluster analysis.

### 1. Introduction

The Rutaceae is a diverse and widely studied family found throughout the world, and its largest genus is *Zanthoxylum*. Many species of *Zanthoxylum* have been extensively studied, due to their large role in the traditional medicine of various cultures<sup>1, 2</sup>. Upon study, these species have demonstrated antifungal, antimicrobial, anti-inflammatory, gastrostimulant, and cytotoxic properties<sup>3-7</sup>. Here, we report the essential oil composition of an undescribed species, *Zanthoxylum* sp. A “Peñas Blancas”.

*Zanthoxylum* sp. A “Peñas Blancas” is a small tree, with glabrous, alternate, parapinnate leaves with 7-8 leaflets. The leaflets, 17×7 cm, are round near the base, grooved dorsally near the apex with petioles around 7-10 cm long; the leaflet apex is abruptly acuminate, the base gradually narrowing and acute, the margins crenate with translucent glands, but the lamina interior without glands. The midvein and lateral veins are impressed above, expressed below, with 8-11 lateral veins per side. The rachis is 10-26 cm long with sparse prickles. The leaves are dark green above and pale olive-green below.

### 2. Materials and Methods

#### 2.1 Plant Material

Leaves of *Zanthoxylum* sp. A “Peñas Blancas” were collected from a non-reproductive individual tree growing in the Peñas Blancas River Valley of the Monteverde Cloud Forest Preserve (10.3091 N, 84.7162 W, 800 m elevation) on May 14, 2008. The plant was characterized by W. A. Haber, and a voucher specimen was deposited in the herbarium of the National Museum of Costa Rica (voucher number Haber ex Cruz 13276). The fresh leaves (45.9 g) were chopped and hydrodistilled for 4 h using a Likens-Nickerson hydrodistillation apparatus with continuous extraction with CHCl<sub>3</sub> (50 mL). The chloroform extract was then evaporated to give the essential oil (618.1 mg).

#### 2.2 Gas Chromatography – Mass Spectrometry

The leaf essential oil of *Zanthoxylum* sp. A “Peñas Blancas” was subjected to gas chromatographic-mass spectral analysis using an Agilent 6890 GC with Agilent 5973

### Correspondence:

**William N. Setzer**

Department of Chemistry, University  
of Alabama in Huntsville  
Huntsville, AL 35899, USA  
Email: [wsetzer@chemistry.uah.edu](mailto:wsetzer@chemistry.uah.edu)

mass selective detector, fused silica capillary column (HP-5ms, 30 m × 0.25 mm), helium carrier gas, 1 mL/min flow rate; injection temperature 200 °C, oven temperature program: 40 °C initial temperature, hold for 10 min; increased at 3 °C/min to 200 °C; increased 2 °C/min to 220 °C, and interface temp 280 °C; EIMS, electron energy, 70 eV. The sample was dissolved in CHCl<sub>3</sub> to give a 1% w/v solution; 1-μL injections using a splitless injection technique were used. Identification of oil components was achieved based on their retention indices (determined with reference to a homologous series of normal alkanes), and by comparison of their mass spectral fragmentation patterns with those reported in the literature<sup>[8]</sup> and stored on the MS library [NIST database (G1036A, revision D.01.00)/ChemStation data system (G1701CA, version C.00.01.08)].

### 2.3 Numerical Cluster Analysis

Forty-four *Zanthoxylum* samples were treated as operational taxonomic units (OTUs). The percentage composition of the main essential oil components was used to determine the chemical relationship between the different samples of *Zanthoxylum* by cluster analysis using the NTSYSpc software, version 2.2<sup>[9]</sup>. Correlation was selected as a measure of similarity, and the unweighted pair-group method with arithmetic average (UPGMA) was used for cluster definition.

### 3. Results and Discussion


The essential oil of the fresh leaves of *Zanthoxylum* “Peñas Blancas” was recovered in 1.35% yield, and a total of 12 components were identified in the oil (Table 1). The oil was composed of 35% monoterpenoid hydrocarbons, 52% oxygenated monoterpenoids, and 13% sesquiterpenoid hydrocarbons. The major components of the oil were citronellal (23.3%), limonene (16.7%), *trans*-pinocampheol (11.6%), and β-pinene (10.8%). Phytochemicals such as citronellal and limonene are found in substantial amounts in many *Zanthoxylum* species<sup>[10-12]</sup>, and many of the lesser abundant components of the oil like α-pinene, linalool, and germacrene D are common throughout the Rutaceae<sup>[13-15]</sup>. The presence of several of the oxygenated monoterpenoids, *trans*-pinocampheol, pinocampheol, myrtenal, and *trans*-dihydrocarvone, however, was unexpected and these components had not been found in any other studied *Zanthoxylum* species.

**Table 1:** Chemical Composition of the leaf oil of *Zanthoxylum* sp. A “Peñas Blancas”.

| RI   | Compound                     | % Composition |
|------|------------------------------|---------------|
| 943  | α-Pinene                     | 7.5           |
| 977  | β-Pinene                     | 10.8          |
| 1025 | Limonene                     | 16.7          |
| 1098 | Linalool                     | 7.3           |
| 1150 | Citronellal                  | 23.3          |
| 1154 | <i>trans</i> -Pinocampheol   | 11.6          |
| 1171 | Pinocampheol                 | 4.9           |
| 1193 | Myrtenal                     | 3.7           |
| 1202 | <i>trans</i> -Dihydrocarvone | 1.3           |
| 1418 | ( <i>E</i> )-Caryophyllene   | 4.3           |
| 1480 | Germacrene D                 | 6.9           |
| 1494 | Bicyclogermacrene            | 1.7           |

In order to place *Zanthoxylum* sp. A “Peñas Blancas” in chemical context with other members of the genus, a cluster analysis of the

major essential oil components has been carried out and is shown in Figure 1. The analysis reveals a great degree of variability and plasticity in *Zanthoxylum* essential oil composition. There is a high degree of intraspecific variation as well as interspecific variation. Thus, for example, the sample of *Z. rhoifolium* from Costa Rica<sup>[13]</sup> is chemically very different from those from Brazil<sup>[16-18]</sup> and *Z. fagara* oils from Cuba<sup>[19]</sup> and Costa Rica<sup>[10]</sup> have very different chemical profiles.



**Fig 1:** Dendrogram obtained by cluster analysis of the percentage composition of essential oils from *Zanthoxylum* samples, based on correlation and using the unweighted pair-group method with arithmetic average (UPGMA).

### 4. Conclusions

The leaf oil composition of *Zanthoxylum* sp. A “Peñas Blancas” as revealed in this study is chemically distinct from other *Zanthoxylum* oils (see above), and this is reflected in the cluster analysis.

### 5. Acknowledgments

We are grateful to the Monteverde Cloud Forest Preserve and the Tropical Science Center for granting us permission to collect plant materials from the Preserve under a cooperative rights agreement and to the Commission for the Development of Biodiversity of Costa Rica’s Ministry of the Environment, Energy, and Telecommunications for Research Permit R-001-2006-OT-CONAGEBIO. We thank Eladio Cruz for collecting leaves of *Zanthoxylum* sp. A “Peñas Blancas”.

## 6. References

1. Matu EN, Staden VJ. Antibacterial and anti-inflammatory activities of some plants used for medicinal purposes in Kenya. *J Ethnopharmacol* 2003; 87:35-41.
2. Arrieta J, Reyes B, Calzada F, Cedillo-Rivera R, Navarrete A. Amoebicidal and giardicidal compounds from the leaves of *Zanthoxylum liebmannianum*. *Fitoterapia* 2001; 72:295-297.
3. Ngane AN, Biyiti L, Zollo PHA, Bouchet PH. Evaluation of antifungal activity of extracts of two Cameroonian Rutaceae: *Zanthoxylum leprieurii* Guill. et Perr. and *Zanthoxylum xanthoxyloides* Waterm. *J. Ethnopharmacol* 2000; 70:335-342.
4. Ngassoum MB, Essia-Ngang JJ, Tatsadjieu LN, Jirovetz L, Buchbauer G, Adjoudji O. Antimicrobial study of essential oils of *Ocimum gratissimum* leaves and *Zanthoxylum xanthoxyloides* fruits from Cameroon. *Fitoterapia* 2003; 74:284-287.
5. Abraham GJS, Agshikar NV. Anti-inflammatory activity of an essential oil from *Zanthoxylum budrunga*. *Pharmacology* 1972; 7:109-114.
6. Itthipanichpong C, Ruangrunsi N, Pattanaautsahakit C. Chemical compositions and pharmacological effects of essential oil from the fruit of *Zanthoxylum limonella*. *J Med Assoc Thai* 2002; 85(suppl:1):S344-S354.
7. Silva DSL, Figueiredo PM, Yano T. Cytotoxic evaluation of essential oil from *Zanthoxylum rhoifolium* Lam. leaves. *Acta Amazonica* 2007; 37:281-286.
8. Adams RP. Identification of Essential Oil Components by Gas Chromatography/Mass Spectrometry, Edn 4, Allured Publishing, Carol Stream, IL, USA, 2007.
9. Rohlf JF. NTSYSpc, Numerical Taxonomy and Multivariate Analysis System. Applied Biostatistics Inc., New York, 2005.
10. Setzer WN, Schmidt JM, Eiter LC, Haber WA. The leaf oil composition of *Zanthoxylum fagara* (L.) Sarg. from Monteverde, Costa Rica, and its biological activities. *J Essent Oil Res* 2005; 17:333-335.
11. Chagonda L, Gundidza M, Wijesuria AMS, Mavi S. Composition of the leaf oil of *Zanthoxylum chalybeum* Thunb. *J Essent Oil Res* 1994; 6:561-563.
12. Wu Y, Shimoda M, Osajima Y. Volatile aroma compounds in young leaves and green fruits of Japanese pepper [*Zanthoxylum piperitum*]. *J Agric Chem Soc Jpn* 1996; 70:1001-1005.
13. Boehme AK, Noletto JA, Haber WA, Setzer WN. Bioactivity and chemical composition of the leaf essential oils of *Zanthoxylum rhoifolium* and *Zanthoxylum setulosum* from Monteverde, Costa Rica. *Nat Prod Res* 2008; 22:31-36.
14. Tiwary M, Naik SN, Tewary DK, Mittal PK, Yadav S. Chemical composition and larvicidal activities of the essential oil of *Zanthoxylum armatum* DC (Rutaceae) against three mosquito vectors. *J Vect Borne Dis* 2007; 44:198-204.
15. Setzer WN, Noletto JA, Lawton RO, Haber WA. Leaf essential oil composition of five *Zanthoxylum* species from Monteverde, Costa Rica *Mol Divers* 2005; 9:3-13.
16. Andrade EHA, Zoghbi MDGB. Constituents of the essential oil of *Zanthoxylum rhoifolium* Lam. *J Ess Oil-Bear Plants* 2004; 7:179-181.
17. Moura DNF, Strapazon JO, Loro F, Morel AF, Flach A. Composition of the leaf oils of Rutaceae: *Zanthoxylum hyemale* A. St. Hill, *Z. rhoifolium* Lam. and *Z. rugosum* A. St. Hill et. Tul. *J Essent Oil Res* 2006; 18:627-628.
18. Silva DSL, Figueiredo PMS, Yano T. Chemotherapeutic potential of the volatile oils from *Zanthoxylum rhoifolium* Lam leaves. *Eur J Pharmacol* 2007; 576:180-188.
19. Pino JA, Agüero J, Marbot R, Fernandes P. Composition of the essential oil of *Zanthoxylum fagara* (L.) Sargent. from Cuba. *J Essent Oil Res* 2005; 17:413-414.
20. Vila R, Iglesias J, Cañigüeral S, Ciccio JF. Composition of the essential oil from leaves of *Zanthoxylum procerum* from Costa Rica. *J Essent Oil Res* 2002; 14:44-46.
21. Guy I, Charles B, Guinaudeau H, Ferreira ME, Rojas de Arias A, Fournet A. Essential oils from *Zanthoxylum chiloperone* and *Z. riedelianum* growing in Paraguay. *Pharmaceut Biol* 2001; 39:152-154.
22. Ivanova A, Kostova I, Navas HR, Villegas J. Volatile components of some Rutaceae species. *Z Naturforsch* 2004; 59c:169-173.
23. Kojima H, Kato A, Kubota K, Kobayashi A. Aroma compounds in the leaves of Japanese pepper (*Zanthoxylum piperium* DC) and their formation from glycosides. *Biosci Biotech Biochem* 1997; 61:491-494.
24. Chagonda L, Gundidza M, Wijesuria AMS, Mavi S. Composition of the leaf oil of *Zanthoxylum chalybeum* Thunb. *J Essent Oil Res* 1994; 6:561-563.
25. Weyerstahl P, Marschall H, Splittgerber U, Son PT, Giang PM, Kaul VK. Constituents of the essential oil from the fruits of *Zanthoxylum rhesoides* Drake from Vietnam and from the aerial parts of *Zanthoxylum alatum* Roxb. from India. *Flavour Fragr J* 1999; 14:225-229.
26. Oyediji AO, Lawal OA, Adeniyi BA, Alaka SA, Tetede E. Essential oil composition of three *Zanthoxylum* species. *J Essent Oil Res* 2008; 20:69-71.
27. Luong NX, Hac LV, Dung NX. Chemical composition of the leaf oil of *Zanthoxylum alatum* Roxb. from Vietnam. *J Essent Oil-Bear Plants* 2003; 6:179-184.
28. Brophy JJ, Goldsack RJ, Forster PI, Hutton I. Composition of the leaf oils of the Australian and Lord Howe Island species of *Zanthoxylum* (Rutaceae). *J Essent Oil Res* 2000; 12:285-291.
29. Feitosa EMA, Arriaga AMC, Lemos TLG, Lima JQ, Nunes e Vasconcelos J, Oliveira MCF, Santiago GMP, Nascimento RF. *Zanthoxylum articulatum* Engler (Rutaceae) essential oil: Chemical composition and larvicidal activity. *J Essent Oil Res* 2007; 19:384-386.
30. Lamaty G, Menut C, Bessiere JM, Aknin M. Aromatic plants of tropical central Africa. II. A comparative study of the volatile constituents of *Zanthoxylum leprieurii* (Guill. et Perr.) Engl. and *Zanthoxylum tessmannii* Engl. leaves and fruit pericarps from the Cameroon. *Flavour Fragr J* 1989; 4:203-205.
31. Joy MT, Verghese J, Retamar JA, Talenti ECJ, Catalán CAN, Gros EG. Essential oil of *Zanthoxylum rhesa*. *Flavour Fragr J* 1986; 1:165-168.